

PREVALENCE OF ORAL CANDIDIASIS IN COMPLETE DENTURES WEARERS

ПРЕВАЛЕНЦА НА ОРАЛНА КАНДИДИЈАЗА КАЈ НОСИТЕЛИ НА ТОТАЛНИ ПРОТЕЗИ

Tmava-Drăgusha A.¹, Shala K.¹, Ivanovski K.², Dirjanska K.², Ristoska S.², Elenchevski S.³

¹Department of Prosthetic Dentistry, Faculty of Medicine, School of Dentistry, Prishtina, Kosovo, ²Department of Oral pathology and Periodontology, Faculty of Dentistry, University "Ss Cyril and Methodius" – Skopje, ³Department of Prosthetic Dentistry, Faculty of Dentistry, University "Ss Cyril and Methodius" – Skopje

Abstract

Objective: As the population age is growing the need for oral prosthetic rehabilitation is increasing accordingly. Oral candidiasis is a common opportunistic infection of the oral cavity caused by an overgrowth of *Candida* spp., the commonest being *Candida albicans*. **Materials and Methods:** To obtain the set objectives, 50 patients (from the experimental group EG), complete denture wearers, were included and sampled at the University Dentistry Clinical Center of Kosovo, Prishtina. Also, we designate a control group of 50 patients, older than 50 years of age. Clinical changes in the oral mucosa were assessed using thorough intra/extra-oral examination with the emphasis on the eventual oral mucosal changes in contact with dentures. In both experimental and control group patients' swabs were obtained for testing the presence of *Candida albicans*. In the experimental group of patients along with oral mucosal swabs, denture swabs were taken as well. **Results:** In our study the highest percentage was found for *Candida albicans* - 26.0%, followed by *Candida tropicalis* - 12.0%, and *Candida crusei* - 6.0%, of the cases. There was a correlation between denture wearing and the occurrence of oral candidiasis. ($p < 0.05$) (Pearson Chi-square: 13.6364, $p = 0.00222$). While there was no correlation found between the duration of denture wearing and the occurrence of oral candidiasis. ($p > 0.05$) (Pearson Chi-square: 1.02339, $p = 0.311716$), however, there was a correlation noted between the years of denture wearing and the occurrence of *Candida* spp. on the patient denture base. **Conclusions:** The prevalence of Denture Stomatitis in complete denture wearers is high and no association with its risk factors was found. **Key words:** denture stomatitis, complete denture wearers, oral candidiasis, *Candida albicans*.

Апстракт

Цел на трудот: Како што се зголемува возраста на популацијата, соодветно се зголемува и потребата од орална протетска рехабилитација. Оралната кандидијаза е честа опортунистичка инфекција во усната празнина, којашто е предизвикана од прекумерен раст на *Candida* spp, од коишто најчесто станува збор за *Candida albicans*. **Материјали и Методи:** За да се постигнат поставените цели, 50 пациенти (од експерименталната група ЕГ), носители на тотални протези, беа вклучени и од нив беа земени примероци во Универзитетскиот Стоматолошки Клинички Центар во Косово, Приштина. Исто, оформивме контролна група од 50 пациенти, постари од 50 години. Клиничките промени на оралната мукоза беа оценети преку темелен интра/екстраорален преглед со акцентирање на евентуалните промени на мукозата којашто е во контакт со тоталната протеза. И во експерименталната и во контролната група, од пациентите беше земен брис за одредување на присуство на *Candida albicans*. Во експерименталната група на пациенти, покрај брисот од оралната мукоза, беше земен и брис од тоталната протеза. **Резултати:** Во нашето испитување, процентуално најголемо присуство беше одредено за *Candida albicans* – 26.0%, потоа *Candida tropicalis* – 12.0% и *Candida crusei* – 6.0% од случаите. Утврдивме корелација помеѓу носењето на тотална протеза и појавата на орална кандидијаза. ($p < 0.05$) (Pearson Chi-square: 13.6364, $p = 0.00222$). Не утврдивме корелација помеѓу времето на носење на протезата и појавата на орална кандидијаза ($p > 0.05$) (Pearson Chi-square: 1.02339, $p = 0.311716$), но, забележавме корелација помеѓу годините на носење на тоталната протеза и појавата на *Candida* spp. на базата на тоталната протеза. **Заклучоци:** Преваленцата на протетскиот стоматитис кај носителите на тотални протези е висока и не е во асоцијација со неговите ризик-фактори. **Клучни зборови:** протетски стоматитис, носители на тотални протези, орална кандидијаза, *Candida albicans*.

Introduction

As the population age is growing the need for oral prosthetic rehabilitation is increasing accordingly¹. Although, the number of treatment options for dentate and edentulous patients is staggering both in number and techniques, not every member of that population can benefit from all of the options, mainly due to the finan-

cial constraints². One of the most common and optimal options, both clinically and financially, is resin-based complete denture³.

Oral fungal infection is a common disease in complete denture wearers. Oral candidiasis is a common opportunistic infection of the oral cavity caused by an overgrowth of *Candida* spp., the most common being *Candida albicans*⁴. *C. albicans* is a dimorphic yeast

strongly gram-positive able to live as a normal commensal organism in the oral cavity of healthy people of 45–65%, but in denture wearers, the prevalence of *Candida* increases to 60–100% which increases the risk of infection^{5,6,7}.

The etiology of denture stomatitis is multifactorial, with factors including trauma caused by ill-fitting dentures, increased age of the patient and of the dentures, lack of denture hygiene, and a favorable environment for proliferation for *Candida albicans* in particular^{8,9,10}.

Patients with candidiasis may report varied symptoms such as painful sensations, local discomfort, difficulty swallowing, a burning sensation in the oral cavity, or an alteration in taste, but such infections are most often asymptomatic¹¹.

Candida species are frequently isolated from the oral cavity in healthy individuals of all ages, and it is therefore difficult to differentiate oral candidiasis from the commensal state by microbiological detection of the *Candida* species in the oral cavity. Therefore, additional microbiological criteria are required to diagnose oral *Candida* infection correctly. Various methods can be used to isolate *Candida* from the oral cavity, including smears, plain swabs, imprint cultures, whole saliva collection, concentrated oral rinses, and mucosal biopsies^{12,13}. Clinical studies have shown that *C. albicans* is not only able to adhere to the mucous surfaces, but also to stick to the acrylic resins of the dental prostheses, where both the plaque accumulated on the dentures and the poor oral hygiene contribute to the virulence of *Candida*, offering a clinical picture of *Candida*-associated denture stomatitis¹⁴.

The purpose of this study was to determine changes in the oral environment occurring in denture-wearers by recording:

- Clinical expression in oral mucosa in close contact with complete dentures.
- Microbiological findings.

Material and Methods

To obtain the set objectives, 50 patients (from the experimental group EG), complete-denture wearers, were included and sampled at the University Dentistry Clinical Center of Kosovo, Department of Prosthodontics, Pristina. Inclusion criteria were as follows:-

- Patients that have worn their dentures for at least one year.
- The dentures should have been fabricated at the laboratory of UDCCK from the same technician under the most similar conditions for denture fabrication for all groups, and

- The patients should be older than 50 years.

Also, we designate a control group (CG) of 50 patients, older than 50 years, without removable prosthodontic restorations, that attend the University Dentistry Clinical Center of Kosovo, Pristina, for other reasons.

History was recorded in the experimental and control group including:

- Medical and Medication History,
- Duration of Denture Wearing (Experimental Group),
- Subjective issues related to their dentures.

Denture hygiene habits were recorded using special questionnaire developed by **Peracini et al.** (2010)¹⁵.

Clinical changes in the oral mucosa were assessed by means of thorough intra/extra-oral examination with emphasis on the eventual oral mucosal changes in contact with dentures. These examinations included the control group as well.

From clinical aspect, all patient with denture related stomatitis are classified into three clinical types (Newton classification):

Newton's type I: pin-point hyperemic lesions (localized simple inflammation)

Newton's type II: diffuse erythema confined to the mucosa contacting the denture (generalized simple inflammation)

Newton's type III: granular surface (inflammatory papillary hyperplasia).

In addition, the dentures' stability was assessed.

In both experimental and control group, patients' swabs were obtained for testing the presence of *Candida albicans*. In the experimental group of patients along with oral mucosal swabs, denture swabs were taken as well.

Microbiology analysis was conducted at the Institute of Public Health, Department of Microbiology, Pristina. Oral and palatal mucosal samples were taken.

Results

In patients from the CG there was no candida infections recorded.

Candida infections were recorded in 24.0% of the patients in the EG. *Candida crusei* and *Candida albicans* was simultaneously found in two patients (4.0%). *Candida albicans* alone was found in 20.0% of patients, evidently *Candida albicans* is found in 24.0% of total patients. (Table 1, Graph 1).

Table 1. Patient distribution by *Candida* – mouth variable


| | CG | | EG | |
|---|-------|---------|-------|---------|
| | Count | Percent | Count | Percent |
| No | 50 | 100.0 | 38 | 76.0 |
| <i>Candida crusei</i> & <i>Candida albicans</i> | | | 2 | 4.0 |
| <i>Candida albicans</i> | | | 10 | 20.0 |
| Total | 50 | 100.0 | 50 | 100.0 |

Table 2. Patient distribution by *Candida*-dentures variable in EG


| | EG | |
|---|-------|---------|
| | Count | Percent |
| No | 35 | 70.0 |
| <i>Candida crusei</i> & <i>Candida albicans</i> | 2 | 4.0 |
| <i>Candida albicans</i> | 7 | 14.0 |
| <i>Candida crusei</i> & <i>Candida tropicalis</i> | 1 | 2.0 |
| <i>Candida tropicalis</i> | 1 | 2.0 |
| <i>Candida albicans</i> & <i>Candida tropicalis</i> | 4 | 8.0 |
| Total | 50 | 100.0 |

In 30% of complete denture wearer patients, candida spp. was isolated on the actual denture base. *Candida crusei* and *Candida albicans* were simultaneously found in two patients (4.0%). *Candida albicans* alone was found in 14.0% of the patients. *Candida crusei* and *Candida tropicalis* were simultaneously found in a single patient, (2.0%). *Candida tropicalis* alone was found in a single patient. *Candida albicans* and *Candida tropicalis* were simultaneously found in 8.0% of EG of patients (Table 2, Graph 2). In total, the highest percentage was found for *Candida albicans* - 26.0%, followed by *Candida tropicalis* - 12.0% and *Candida crusei* - 6.0%, of the cases.

There was a correlation between denture wearing and the occurrence of oral candidiasis. ($p < 0.05$) (Pearson Chi-square: 13.6364, $p = .000222$).



Graph 1. Patient distribution by *Candida* – mouth variable



Graph 2. Patient distribution by *Candida*-dentures variable in EG

During the study, there was no correlation found between the years of denture wearing and the occurrence of oral candidiasis. ($p > 0.05$) (Pearson Chi-square: 1.02339, $p = .311716$), however, there was a correlation noted between the years of denture wearing and the occurrence of *Candida* spp. on the patient denture base. ($p < 0.05$) (Pearson Chi-square: 13.7188, $p = .000212$). Similarly, a correlation between diabetes in denture-wearing patients and *Candida* infections was confirmed. ($p < 0.05$) (Pearson Chi-square: 17.9107, $p = .000023$).

Discussion

The *Candida* species form the normal commensal fungi component, and its activity is regulated by many intrinsic and extrinsic factors. The pathogenic nature of *Candida* has been correlated with various systemic conditions, which in some manner affect the immunity of the patients¹⁶.

The *Candida* concentration associated with several clinical oral signs in the infected patients and may be closely related to the patient's current clinical status and prognosis¹⁷. In our research of patients from the CG there was no *Candida* infections recorded. Our results agree with Al Tarawneh et al.¹⁸ who concluded that mucosal candidal counts and the presence of cytological hyphae did not show significant difference when comparing DS to healthy participants. Our results show that the highest percentage was found for *Candida albicans* in denture in EG patients (26.0%), followed by *Candida tropicalis* and *Candida crusei*, of the cases. Many authors have reported that *Candida albicans* is the most common species in denture users, making up 70% of all isolates¹⁹. Zomordian et

al.²⁰, investigated risk factors associated with progression to *Candida*-related dentures stomatitis in patients using complete dentures, and they found that *C. albicans* was the most frequently recovered species, followed by *C. glabrata* and *C. tropicalis*. Therefore, according to Mizugai et al.²¹ in their study evaluate the association among age distribution and denture wearing status and *Candida* spp. detection rate. This study indicates that, although detection rate of single *Candida* spp. were rather high in the youngest group of non-dentures wearers, detection rates of single and multiple *Candida* spp. were significantly higher in denture wearers of all other age groups compared with non-denture wearers ($P < 0,05$). Many studies have also shown that denture hygiene practices are essential, because the porous material of the surface biofilm can serve as a reservoir of fungal microorganisms, and contribute to reinfection^{22,23}.

During the study there was no correlation found between the years of denture wearing and occurrence of oral candidiasis, however, there was a correlation between the years of denture wearing and occurrence of *Candida* spp. on the patient denture base. Our results agree with previous reports which indicate that DRS are in statistically significant continuous denture wearing^{24,25}. Due to the deterioration of dentures over time, such as the polished surfaces fit to the underlying tissues and the occlusion, dentures could become more irritant to the mucosa and more open to *Candida* and bacterial colonization.

Our findings suggest that there is a significant difference between diabetes in denture wearing patients and candidiasis. The most frequent oral signs and symptoms observed in both controlled and uncontrolled diabetic

patients was hyposalivation, halitosis, periodontitis, taste alteration, aphthous stomatitis, and they found significant difference between the groups. Also, the most common lesion seen in both the groups were candidiasis, fibrous hyperplasia, aphthous ulcers and benign neoplasia where the chi-square test showed a significant difference between the groups $P < 0,05$ according to Shrimali et al.²⁶, Daniluk et al.²⁷. *Candida albicans*, statistically significantly, more frequently was isolated in denture wearers' patients with diabetes melitus and without diabetes, comparing to such groups of patients but without dentures.

Our results may differ due to our smaller number of participants, or due to other issues associated with their age and general health. Therefore it should be pointed out that older dentures are harder to clean due to a tendency for porosity in their bases, which can contribute for the emergence of the disease. It should be concentrated on patient information and motivation on hygiene, for prophylactic purposes. Future studies will be a series of clinical epidemiologic studies, already planned in our department, to investigate pathogenesis of Denture stomatitis.

Denture Wearers

Laura Lourenço Morel¹, Anna Paula da FernandaFaot¹, Luciana de Rezende Pinto¹,

Laura Lourenço Morel¹, Anna Paula da FernandaFaot¹, Luciana de Rezende Pinto¹

Conclusion

In conclusion, the prevalence of Denture Stomatitis in the total number of denture wearers is high and although no association with its risk factors was found.

Preventive and educational measures for oral health in the elderly, stimulating the correct hygiene of the denture and the oral cavity, should be a routine among professionals. Patients that use dentures must maintain their dentures in order to preserve their oral health.

Conflicts of interest

The authors confirm that this article content has no conflict of interest.

Reference

- Ageing WP. World Population Ageing 1950-2050. Popul English Ed [Internet]. 2002;26:5–9. Available from: <http://www.un.org/esa/population/publications/worldageing1950050/>
- Leles CR, Ferreira NP, Vieira AH, Campos AC V, Silva ET. Factors influencing edentulous patients' preferences for prosthodontic treatment. J. Oral Rehabil. 2011;38:333–9.
- McKenna G, Allen F, Woods N, O'Mahony D, Cronin M, Damata C, Normand C. Cost-effectiveness of tooth replacement strategies for partially dentate elderly: a randomized controlled clinical trial. Community Dent. Oral Epidemiol. [Internet]. 2013;1–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2425138>
- Salerno C, Pascale M, Contaldo M, Esposito V, Petrucci M, Sercipo R. Candida-associated denture stomatitis. Med. Oral. Patol. Oral Cir. Bucal. 2011 Mar 1;16 (1):139-43.
- Loster BW, Loster J, Wiecezorek A, Ryniewicz W. Mycological analysis of the oral cavity of patients using acrylic removable dentures. Gastroenterol. Res. Pract. 2012. 2012: 951572. Published online 2012 Apr 8. doi: 10.1155/2012/951572.
- Loster JE, Wiecezorek A, Loster BW: correlation between age and gender in Candida species infections of complete denture wearers: a retrospective analysis. ClinInterv Aging. 2016;11:1707–1714.
- Zomorodian K, Haghighi NN, Rajaei N, et al. Assessment of Candida species colonization and denture-related stomatitis in complete denture wearers. Med. Mycol. 2011;49:208–211.
- Abaci O, Haliki-Uztan A. Investigation of the susceptibility of Candida species isolated from denture wearers to different antifungal antibiotics. Afr. J. Microbiol. Res. 2011;5:1398–1403.
- Wilson J. The aetiology, diagnosis and management of denture stomatitis. Br. Dent. J. 1998;185:380–384.
- Webb BC, Thomas CJ, Willcox MDP, et al. Candida-associated denture stomatitis: aetiology and management: a review: part I: factors influencing distribution of Candida species in the oral cavity. Aust. Dent. J. 1998;43:45–50.
- Hoshi N, Mori H, Taguchi H, et al. Management of oral candidiasis in denture wearers. J. Prosthodont. Res. 2011;55:48–52.
- Williams DW, Lewis MA. Isolation and identification of Candida from the oral cavity. Oral Dis. 2000;6:3-11.
- Byadarahally.Raju S, Rajappa S. Isolation and identification of candida from the oral cavity. ISRN Dent. 2011;2011:487921.doi:10,5402/2011/487921.
- Akpan A, Morgan R. Oral candidiasis. Postgrad. Med. J. 2002;78:455-459.
- Peracini A, de Andrade I. M., da Silva C. H. L., Paranhos H. de F. O. Behaviors and Hygiene Habits of Complete Denture Wearers. Braz. Dent. J. 2010;21(3): 247-252.
- Fenn SM, Narayanan M, Jacob M. Prevalence of oral Candida in saliva of uncontrolled and controlled type 2 diabetes mellitus patients – Beyond reasonable doubt? SRM J. Res..Dent. Sci.2019;10:1-6.
- Tooyama H, Matsumoto T, Hayashi K, Kurashina K, Kurita H, Uchida M, Kasuga E, Honda T. Candida concentrations determined following concentrated oral rinse culture reflect clinical oral signs. BMC Oral Health 2015; 15:150.
- Al Tarawneh S, Bencharit S, Mendoza L, Curran A, Barrow D, Barros S, Preisser , Loewy ZG, Gendreau L, Offenbacher S. Clinical and histological findings of denture stomatitis as related to intraoral colonization patterns of Candida albicans, salivary flow, and dry mouth. J. Prosthodont. 2013 Jan;22(1):13-22.
- D'aistan S, Aktas AE, Caglayan F, Ayyildiz A, Bilge M. Differential diagnosis of dentur-induced stomatitis, Candida, and their variations in patients using complete denture: a clinical and mycological study. Mycoses. 2009;52:266–271. doi:10.1111/j.1439-0507.2008.01592.
- Zomorodian K, Haghighi NN, Rajaei N, Pakshir K, Tarazooie B, Vojdani M, Sedaghat F, Vosoghi M. Assessment of Candida species colonization and denture-related stomatitis in complete denture wearers. Med. Mycol. 2011 Feb;49(2):208-11. doi: 10.3109/13693786.2010.507605. Epub 2010 Aug 26. PubMed PMID: 20795762.
- Mizugai, Hiroyuki; Isogai, Emiko; Hirose, Kimiharu; Chiba, Itsuo. 2007 Effect of denture wearing on occurrence of Candida

-
- species in the oral cavity. *The Journal of Applied Research*. 2007; Vol. 7, No. 3: 250-254.
22. Hoshing C, Dixit S, Mootha A, Diwan N. Role of *Candida albicans* in denture stomatitis. *J. Indian Acad. Oral Med. Radiol*. 2011;23:617–619.
23. Dantas APFM, Consani RLX, Sardi JCO, Mesquita MF, Silva MCVS, Sinhoreti MAC: biofilm formation in denture base acrylic resins and disinfection method using microwave. *J. Res. Pract. Dent*. 2014; Article ID 112424. doi:10.5171/2014.112424.
24. Fenlon MR, Sherriff M, Walter JD. Factors associated with the presence of denture related stomatitis in complete denture wearers: a preliminary investigation. *Eur. J. Prosthodont. Restor. Dent*. 1998; 6:145-147.
25. Jeganathan S, Payne JA, Thean HP. Denture stomatitis in an elderly edentulous Asian population. *J. Oral Rehabil*. 1997; 24:468-472.
26. Shrimali L, Astekar M, Sowmya GV. Correlation of Oral Manifestations in Controlled and Uncontrolled Diabetes Mellitus. *International Journal of Oral & Maxillofacial Pathology*. 2011;2(4):24-27.
27. Daniluk T, Tokajuk G, Stokowska W, Fiedoruk K, Sciepek M, Zaremba ML, Rozkiewicz D, Cylwik-Rokicka D, Kedra BA, Anielska I, Górska M, Kedra BR. Occurrence rate of oral *Candida albicans* in denture wearer patients. *Adv. Med. Sci*. 2006;51Suppl 1:77-80.