

THE DISTRIBUTION OF OCCLUSAL VERTICAL STRESS IN SHORTENED DENTAL ARCHS WITH CROSS-ARCH DENTAL BRIDGES

ДИСТРИБУЦИЈА НА ОКЛУЗАЛНИ ВЕРТИКАЛНИ СИЛИ КАЈ МАЛКУ СКРАТЕНИ ЗАБНИ НИЗИ СО МОСТОВНИ КОНСТРУКЦИИ

Vujasin S.¹, Bundevska J.², Kokalanov V.³, Dejanoska T.⁴, Vankovski V.⁵

¹Faculty of Dentistry, Department of Prosthodontics, EURM, Skopje, RM, ²Faculty of Dentistry, Department of Prosthodontics, UKIM, Skopje, RM, ³Faculty of Computer Science, Numerical Analysis and Applied Mathematics, UGD, Shtip, RM, ⁴PZU Dental International, Skopje, ⁵Faculty of Dentistry, Department of Prosthodontics, EURM, Skopje,

Abstract

The **aim** of the research is to analyze the distribution of occlusal vertical forces of the abutment teeth in bridge constructions with slightly shortened dental arches. **Material and method.** Subject of research are bridge constructions in slightly shortened dental arches in the mandibula. The Finite Element Method (FEM) was applied, and the analysis was made with simultaneous both-sided loading on the distal three abutment teeth. The simulated loading forces are with strength between 0.5 to 512N. Distributed occlusal vertical forces are measured at the level of the periodontal ligament (PDL). **Results.** The results show a symmetrical distribution of forces on both sides of the bridge structure. The largest forces are distributed on teeth with applied force. The smallest forces are distributed in the apical part of PDL. There was no difference between the total distributed forces on teeth and the distributed forces on the lateral part of the PDL. The measured strength of the forces and the percentage are within the known values in the literature. **Conclusion.** The obtained results of the research are within the level of known functional values of the mastication forces. This means that the biomechanical aspect for bridge constructions with slightly shortened dental arches is not disputable, but the decision to make it should also be based on other individual and clinical conditions. **Keywords:** Occlusal vertical forces, bridge construction, shortened dental arches, finite elements method (FEM), periodontal ligament (PDL).

Апстракт

Цел на трудот е да се направи анализа на дистрибуцијата на оклузални вертикални сили на забите носачи кај мостовни конструкции кај малку скратени забни низи. **Материјал и метод.** Предмет на истражување е мостовна конструкција кај малку скратен забен низ во долна вилица. Применет е метод на конечни елементи (МКЕ) а анализата е направена со симетрично оптоварување на дисталните три заби носачи. Симулираните сили на оптоварување се со јачина од 0.5 до 512N. Дистрибуираните сили се мерени на ниво на перодонталниот лигамент (ПДЛ). **Резултати.** Резултатите покажуваат симетрична дистрибуција на силите и на двете страни на мостовната конструкција. Најголеми сили се дистрибуираат на забите на кои делува аплицираната сила. Најмали сили се дистрибуираат во апикалниот дел на ПДЛ. Не е најдена разлика помеѓу вкупно дистрибуираните сили на забот и дистрибуираните сили на страничниот дел на ПДЛ. Измерените сили по јачина и процент се во рамките на познатите вредности во литературата. **Заклучок.** Добиените резултатите од истражувањето се во ниво на познатите функционални вредности на џвакалните сили. Ова значи дека од биомеханички аспект изработка на мостовни конструкции кај малку скратени забни низи не е спорна, но одлуката за изработка треба да се донесе и врз база на другите индивидуални и клинички услови. **Клучни зборови:** Оклузални вертикални сили, мостовни конструкции, скратени забни низи, метод на конечни елементи (МКЕ), периодонтален лигамент (ПДЛ).

Introduction

Until the seventies of the twentieth century, the goal of dental treatment was the maintenance of complete dental arches with 28 teeth. In 1981, the concept of the shortened dental arch (SDA) promoted by Käyser, suggesting that shortened dental arches with at least four occlusal units, preferably in a symmetrical position, have sufficient capacity to maintain an adequate oral function¹.

This claim caused a series of researches in which the concept of shortened dental arch was studied from all aspects, the functions of the masticatory system, the effect of dental treatment, the quality of life, as well as the economic and social aspects^{2, 3, 4, 5, 6}.

Based on the results of extensive surveys, the World Health Organization WHO in 1992 set the new goal of dental treatment, which is a healthy, natural, and function-

al dental arch with at least 20 teeth without the need for prosthesis^{7,8}.

Anneloes and associates found that shortened dental arches can stay stable for more than 27 years, which justifies the dental concept for shortened dental arches⁹.

However, certain teeth changes can disrupt the stability of the shortened dental arches and in that case, a prosthetic treatment would be needed. The most frequent changes are the migration of teeth by separation, aesthetical and functional needs, and especially the loss of bone support.

According to De Oliveira and associates, teeth in shortened arches show greater movement than teeth in intact dental arches¹⁰.

According to Kourkout and associates, bridge constructions can provide a certain degree of rigidity and enable a more favorable distribution of the masticatory forces of all remaining cases in patients with an advanced degree of progressive changes in the periodont¹¹.

Measuring the the masticatory forces is important in order to assess the functional state of the masticatory system, but it is a complex problem. The values obtained depend on many factors in the masticatory system, in the body, as well as of the measurement methods. Therefore, we find differences in the values obtained in literature.

In the published results, prevailing data suggest that the occlusal forces in static occlusion range from 100 to 1000N, while the data of the functional masticatory forces range from 3.5 to 350N¹².

According to Waltimo and Kononen, the maximal occlusal forces between Europeans and Americas range from 600 to 750N, while the functional mastication forces are much smaller, around 60-100 N¹³.

Veski measured the maximal masticatory forces in intact dental arches in the lower jaw from 176.8 N to 380.9 N in women and from 193.7 N to 506.9 N in men¹⁴.

Himmlová measured the approximate value of 135 N in masticatory forces in subjects with natural intact dental arches¹⁵.

Laurell L. measured maximal masticatory forces of 320 N in circular dental bridge constructions with a healthy periodont and 264 N with weakened paradont¹⁶.

In his research, Biswas found the mean values of the maximal masticatory forces for the front teeth to be 193N, for the canines 223 N, for the premolars 280 N and for the molars 350N¹⁷.

The mean value of the occlusal force is between 39-66 N for the premolars and 11-33 N for the front teeth¹⁸.

According to Lundgren and Laurell, the maximal force that occurs during the chewing act is 280N and the medium functional force is about 100N. He thinks that, on average, about 37% of the total maximal occlusal force is used in chewing¹⁹.

Sato too believes that the dynamic functional momentum is 35-45% of the measured static forces²⁰.

By electromyographic analysis of the masticatory muscles, Prochechel and Morneburg found a mean functional masticatory force of 220N²¹.

Several authors noted that the strength of the masticatory forces increases in the distal direction of dental arches^{16, 22, 23, 24, 25, 26, 27, 28}.

Kondo measured higher values of the maximal masticatory force in slightly shortened dental arches and pressure on PDL on second premolars²⁹.

Guo and associates and André and associates proved that the deformation forces of the periodontal ligament increase in proportion to the increase of the loading^{30,31}.

According to Apostolov, there are no distinctive differences in the values of the maximal masticatory forces on the left and on the right side of the dental arches^{32,33}.

Cai and associates and Zhou Shu-min and associates found out that in the initial teeth loading, a larger amount of the masticatory forces are distributed on the cervical and lateral parts of the periodontal ligament^{34,35,36}.

In Jayam's research, during vestibular, lingual and incisal loading, the distribution of force in the apical zone is from 0.75 to 0.80N³⁷.

Fratila, Oruč and Je J discovered that the greatest concentration of the forces is distributed at the site of force action^{18, 38, 39}.

Al-Zarea measured higher values of force on the side of the natural teeth in comparison to the bridge structure side, in same subjects, the difference was statistically significant ($p < 0.05$)⁴⁰.

According to literature data, the highest influence on teeth is inflicted by the occlusal vertical forces which are the largest and the base of mastication. They are one of the most important conditions for physiologically optimal occlusion, especially in prosthetic constructions where the tendency during modeling is to reduce the impact of side forces²⁴.

From this aspect, the interest of this research is the distribution of the occlusal vertical forces of bridge constructions made in small shortened sequences.

Aim of the research

The aim of this research is to analyze the distribution of occlusal vertical dental forces of bridge constructions in slightly shortened dental arches.

Material and methods

The analysis of the distribution of occlusal vertical forces is made by using the finite element method. The research was done on a three-dimensional computer

model, the computer analysis and the generation of the finite element network were done with SOFISTIK software package.

The basic model is Kennedy class I edentulous in the mandibula with end teeth 45 and 35, a slightly shortened dental arch (Figure 1.).




Figure 1. Model of bridge construction in slightly shortened dental arch

The discretization of the model is on finite elements with six sides and eight nodes.

The values needed for modeling teeth, periodontal ligament, component materials for bridge structures were taken from literature data.

The analysis was performed by symmetrical loading on the three distal abutment teeth of the bridge construction 45, 44, 43, 42, 41, 31, 32, 33, 34, 35 (Figure 2).




Figure 2. Symmetrical loading of the three distal teeth

Simulated vertical occlusal forces with a magnitude of 0.5N to 512N were applied in the research, respectively to the literature data for maximal and minimal masticatory forces.

The duration of the load force is not taken as a factor for the analysis.

The applied finite element method is a recognized and used method in dental researches, especially in dental biomechanics.

In this research, a three-dimensional (3D) finite element method (FME) is applied, based on the deformation method.

It is a numerical method that performs physical discretization of space. The continuum (independent of its shape, size) is divided into elements with finite dimensions. These elements are connected to each other in discrete points marked as nodes, and that way, a finite element network is formed. By analyzing the finite elements, actually the analysis of the continuum as a whole is carried out.

The data for teeth, periodontal ligament, materials used for bridge structures are taken from literature^{41,42}.

The problem being analyzed in this research is nonlinear. Nonlinearity is due to the anisotropic properties of the periodontal ligament. According to Kojima, PDL should be modeled as nonlinear⁴³.

By using the nonlinear analysis, a real response is obtained for the behavior of bridge structures and the distribution of forces.

Results

By carrying out the foreseen examinations in the research, we obtained the following results:

Table 1 shows the obtained values of the distributed occlusal vertical forces on the abutment teeth during simultaneous both-sided load of the three distal abutment teeth 45 44 43 and 33 34 35 in bridge construction at 45 44 43 42 41 31 32 33 34 35 with the slightly shortened dental arch.

Table 2 shows the percentage distribution of occlusal vertical forces on the abutment teeth during simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35 in bridge construction 45 44 43 42 41 31 32 33 34 35 with the slightly shortened dental arch.

Table 3 shows the obtained values of the distributed occlusal vertical forces in the apical part of PDL of abutment teeth during simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35 in bridge construction 45 44 43 42 41 31 32 33 34 35 with a slightly shortened dental arch.

Table 4 shows the percentage distribution of occlusal vertical forces in the apical part of PDL of abutment teeth during simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35 in bridge construction 45 44 43 42 41 31 32 33 34 35 with a slightly shortened dental arch.

Table 1. Values of the distributed occlusal vertical forces on the abutment teeth during simultaneous both-sided load of the three distal abutment teeth 45 44 43 and 33 34 35

F Tooth	0.5 N	1 N	2 N	4 N	8 N	16 N	32 N	64 N	128 N	256 N	512 N
45	-0.17	-0.35	-0.69	-1.39	-2.92	-5.89	-13.58	-24.85	-49.91	-94.30	-186.09
44	-0.13	-0.26	-0.52	-1.05	-1.96	-4.24	-7.20	-18.52	-35.54	-79.09	-160.03
43	-0.10	-0.20	-0.40	-0.80	-1.56	-2.97	-5.53	-11.27	-25.92	-52.13	-113.78
42	-0.05	-0.11	-0.21	-0.42	-0.84	-1.55	-3.05	-4.74	-10.24	-18.44	-31.18
41	-0.04	-0.08	-0.17	-0.34	-0.70	-1.32	-2.56	-4.09	-6.78	-11.80	-16.31
31	-0.04	-0.08	-0.17	-0.34	-0.69	-1.32	-2.56	-4.09	-6.74	-11.76	-16.28
32	-0.05	-0.11	-0.21	-0.42	-0.84	-1.54	-3.06	-4.74	-10.28	-18.49	-31.33
33	-0.10	-0.20	-0.40	-0.80	-1.56	-2.96	-5.54	-11.18	-25.86	-52.02	-113.56
34	-0.13	-0.26	-0.52	-1.05	-1.96	-4.24	-7.21	-18.49	-35.50	-79.11	-160.04
35	-0.17	-0.35	-0.70	1.39	-2.92	-5.89	-13.60	-24.87	-49.94	-94.37	-186.18
Total	-1.00	-2.00	-3.99	-7.99	-15.98	-31.94	-63.88	-126.83	-256.71	-511.51	-1014.78

Table 2. Percentage distribution of occlusal vertical forces on the abutment teeth during simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35

F Tooth	0.5 N	1 N	2 N	4 N	8 N	16 N	32 N	64 N	128 N	256 N	512 N
45	17.40	17.36	17.40	17.37	18.30	18.44	21.26	19.59	19.44	18.44	18.45
44	13.12	13.10	13.12	13.10	12.29	13.29	11.27	14.660	13.84	15.46	15.77
43	10.04	10.04	10.03	10.03	9.79	9.31	8.66	8.88	10.10	10.19	11.21
42	5.25	5.27	5.24	5.26	5.27	4.84	4.78	3.74	3.99	3.61	3.07
41	4.20	4.24	4.21	4.24	4.35	4.15	4.01	3.22	2.64	2.31	1.61
31	4.20	4.24	4.20	4.23	4.35	4.14	4.00	3.22	2.63	2.30	1.60
32	5.24	5.26	5.24	5.25	5.27	4.84	4.79	3.74	4.01	3.61	3.09
33	10.03	10.02	10.03	10.02	9.78	9.27	8.67	8.82	10.07	10.17	11.19
34	13.12	13.09	13.13	13.10	12.30	13.27	11.28	14.58	13.83	15.47	15.77
35	17.42	17.37	17.42	17.38	18.30	18.45	21.29	19.61	19.45	18.45	18.35
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

Table 3. Values of the distributed occlusal vertical forces in the apical part of PDL

F Tooth	0.5 N	1 N	2 N	4 N	8 N	16 N	32 N	64 N	128 N	256 N	512 N
45	-0.001	-0.002	-0.003	-0.006	-0.013	-0.026	-0.079	-0.151	-0.270	-0.443	-0.817
44	-0.001	-0.001	-0.002	-0.005	-0.009	-0.019	-0.031	-0.110	-0.205	-0.383	-0.704
43	0.000	-0.001	-0.001	-0.003	-0.006	-0.011	-0.031	-0.048	-0.126	-0.230	-0.429
42	0.000	0.000	-0.001	-0.002	-0.003	-0.006	-0.011	-0.018	-0.047	-0.091	-0.139
41	0.000	0.000	-0.001	-0.001	-0.003	-0.005	-0.009	-0.015	-0.020	-0.053	-0.074
31	0.000	0.000	-0.001	-0.001	-0.003	-0.005	-0.009	-0.015	-0.020	-0.053	-0.074
32	0.000	0.000	-0.001	-0.002	-0.003	-0.006	-0.012	-0.019	-0.049	-0.095	-0.145
33	0.000	-0.001	-0.001	-0.003	-0.006	-0.011	-0.020	-0.047	-0.125	-0.228	-0.425
34	-0.001	-0.001	-0.002	-0.005	-0.009	-0.019	-0.031	-0.110	-0.205	-0.383	-0.704
35	-0.001	-0.002	-0.003	-0.006	-0.013	-0.027	-0.079	-0.151	-0.270	-0.443	-0.818
Total	0.00	-0.01	-0.02	-0.03	-0.07	-0.13	-0.30	-0.68	-1.34	-2.40	-4.33

Table 4. Percentage distribution of occlusal vertical forces in the apical part of PDL on abutment teeth

F Tooth	0.5 N	1 N	2 N	4 N	8 N	16 N	32 N	64 N	128 N	256 N	512 N
45	0.08	0.08	0.08	0.08	0.08	0.08	0.12	0.12	0.11	0.09	0.08
44	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.09	0.08	0.07	0.07
43	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.04	0.05	0.04	0.04
42	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.02	0.02	0.01
41	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01
31	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01
32	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01
33	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.04	0.05	0.04	0.04
34	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.09	0.08	0.07	0.07
35	0.08	0.08	0.08	0.08	0.08	0.08	0.12	0.12	0.11	0.09	0.08
Total	0.42	0.42	0.42	0.42	0.42	0.42	0.47	0.54	0.52	0.47	0.43

Table 5. Values of distributed occlusal vertical forces on the lateral parts of PDL of abutment teeth during simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35

Tooth \ F	0.5 N	1 N	2 N	4 N	8 N	16 N	32 N	64 N	128 N	256 N	512 N
45	-0.17	-0.35	-0.69	-1.38	-2.91	-5.86	-13.50	-24.70	-49.64	-93.86	-185.27
44	-0.13	-0.26	-0.52	-1.04	-1.96	-4.22	-7.17	-18.41	-35.33	-78.70	-159.32
43	-0.10	-0.20	-0.40	-0.80	-1.56	-2.96	-5.51	-11.22	-25.80	-51.90	-113.35
42	-0.05	-0.10	-0.21	-0.42	-0.84	-1.54	-3.04	-4.73	-10.20	-18.35	-31.04
41	-0.04	-0.08	-0.17	-0.34	-0.69	-1.32	-2.55	-4.07	-6.76	-11.74	-16.23
31	-0.04	-0.08	-0.17	-0.34	-0.69	-1.32	-2.55	-4.07	-6.72	-11.71	-16.20
32	-0.05	-0.10	-0.21	-0.42	-0.84	-1.54	-3.05	-4.72	-10.23	-18.39	-31.18
33	-0.10	-0.20	-0.40	-0.80	-1.56	-2.95	-5.52	-11.13	-25.73	-51.80	-113.14
34	-0.13	-0.26	-0.52	-1.04	-1.96	-4.22	-7.18	-18.38	-35.29	-78.73	-159.33
35	-0.17	-0.35	-0.69	-1.38	-2.91	-5.87	-13.53	-24.72	-49.67	-93.93	-185.36
Total	-1.00	-1.99	-3.98	-7.95	-15.91	-31.80	-63.58	-126.15	-255.38	-509.11	-1010.45

Table 6. Percentage distribution of occlusal vertical forces on lateral parts of PDL in abutment teeth

Tooth \ F	0.5 N	1 N	2 N	4 N	8 N	16 N	32 N	64 N	128 N	256 N	512 N
45	17.32	17.28	17.32	17.30	18.21	18.36	21.14	19.47	19.34	18.35	18.26
44	13.06	13.04	13.06	13.04	12.24	13.23	11.22	14.51	13.76	15.39	15.70
43	10.00	10.01	9.99	10.00	9.75	9.27	8.63	8.85	10.05	10.15	11.17
42	5.23	5.25	5.22	5.24	5.26	4.82	4.76	3.73	3.97	3.59	3.06
41	4.18	4.23	4.19	4.22	4.34	4.13	3.99	3.21	2.63	2.30	1.60
31	4.18	4.22	4.18	4.22	4.33	4.12	3.99	3.21	2.62	2.29	1.60
32	5.22	5.24	5.22	5.23	5.25	4.82	4.77	3.72	3.99	3.60	3.07
33	9.99	9.98	9.99	9.98	9.74	9.24	8.63	8.78	10.02	10.13	11.15
34	13.06	13.03	13.07	13.04	12.24	13.21	11.23	14.49	13.75	15.39	15.70
35	17.34	17.29	17.34	17.31	18.22	18.37	21.17	19.49	19.35	18.36	18.27
Вкупно	99.58	99.58	99.58	99.58	99.58	99.58	99.53	99.46	99.48	99.53	99.57

Table 5 shows the obtained values of the distributed occlusal vertical forces on the side parts of the PDL on abutment teeth during simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35 in bridge construction 45 44 43 42 41 31 32 33 34 35 with a slightly shortened dental arch.

Table 6 shows the percentage distribution of occlusal vertical forces on lateral parts of PDL of abutment teeth during the simultaneous both-sided load on the three distal abutment teeth 45 44 43 and 33 34 35 in bridge construction 45 44 43 42 41 31 32 33 34 35 with a slightly shortened dental arch.

Discussion

The obtained results for the distribution of the occlusal vertical forces on the abutment teeth of bridge constructions in slightly shortened dental arches during simultaneous both-sided loading of the three distal abutment teeth 45 44 43 and 33 34 35 (Table 1) have an approximately identical distribution of the applied force to the left and right side. Such results were also obtained by Apostolov³².

All distributed forces have the same direction of action identical to the direction of action of the loading force.

The distributed force has a tendency to approximately double the rise with the increase in the strength of the applied force. That is in line with the results of Guo and associates and André and associates^{30, 31}.

For all applied loading forces, the strongest force is distributed to the distal abutment teeth that gradually decrease mesially. Fratila, Oruc and Ye Y. found this kind of force distribution^{18, 38, 39}.

On teeth not exposed to direct loading forces, smaller forces are distributed that reach values of 31.33 N for the strongest applied force (Table 1).

During the simultaneous both-sided load of the three distal abutment teeth 45, 44, 43 and 33, 34, 35 in bridge structures, there is approximately identical distribution percentage of distributed force on the left and right side. The percentage of distributed forces is approximately identical for small and large forces with a tendency of slight increase by increasing the strength of the applied force. The highest percentage of force is distributed to distal abutment teeth that gradually decrease to the mesial abutment teeth.

On teeth with no force applied, the greatest force is distributed during small loading forces and gradually decreases with the rise of loading force (Table 2).

The strength of the distributed forces on the apical part of the PDL on the abutment teeth is minimal, and is less than 1 N, which is in accordance with Jayam's research.³⁷

The percentage of the distributed forces of the apical parts of the PDL of the teeth carriers is less than 0.5% (Table 4).

Distributed force on the side parts of the PDL on the abutment teeth has the same characteristics as the distributed forces on teeth. This means that the major part of the force is received by the lateral parts of PDL. This is consistent with the results of Cai and Associates and Zhou Shu-min and associates^{34, 35, 36}.

Over 99.5% of the applied force is distributed to the lateral parts of PDL of the abutment teeth and has the same characteristics as on the entire teeth (Table 6).

The greatest difficulty in this research was that there are many data in the literature, but the results are difficult to compare because different research methods have been used. The published clinical trials are often reduced to periodic analyzes.

Conclusion

Distributed occlusal vertical forces on all abutment teeth have the same direction of action that is identical with the direction of action of the loading force.

For all applied loading forces, the strongest force is distributed to the distal teeth.

The percentage of distributed occlusal vertical forces on teeth is approximately identical for small and large forces.

Distributed occlusal vertical forces on the lateral parts of PDL have the same characteristics as the distributed forces on teeth.

The strength and percentage of the distributed occlusal vertical forces on the apical part of PDL on the abutment teeth is minimal, and is less than 1N or 0.5%.

Reference

1. Käyser AF. Shortened dental arches and oral function. *J Oral Rehabil.* Sep;8(5):457-62.
2. Aras K., Hasanreisoglu U., Shinogaya T. Masticatory performance, maximum occlusal force, and occlusal contact area in patients with bilaterally missing molars and distal extension removable partial dentures. *Int J Prosthodont.* 2009; 22(2): 204-9.
3. Paulo Tabunyangi Sarit. The shortened dental arch concept and its relevance for oral health care in developing countries. *International Journal of Contemporary Dentistry*, Vol 3, No 1 (2012)
4. S, Musekiwa A, Chikte UME, Omar R (2014). Differences in Functional Outcomes for Adult Patients with Prosthodontically-Treated and - Untreated Shortened Dental Arches: A Systematic Review. *PLoS ONE* 9(7): e101143. <https://doi.org/10.1371/journal.pone.0101143>

5. Wolfart S. The randomized shortened dental arch study: oral health-related quality of life. *Clinical Oral Investigations* March 2014; 18 (20): 525-533.
6. Fueki K1, Igarashi Y1, Maeda Y2, Baba K3, Koyano K4, Sasaki K5, Akagawa Y6, Kuboki T7, Kasugai S8, Garrett NR9. Effect of prosthetic restoration on masticatory function in patients with shortened dental arches: a multicentre study. *J Oral Rehabil.* 2016 Jul;43(7):534-42. doi: 10.1111/joor.12387. Epub 2016 Feb 8.
7. Jepsen NJ, Allen PF. Short and sticky options in the treatment of the partially dentate patient. *Br Dent J* 1999;187:646-52.
8. Marcus S.E., Drury T.F., Brown L.J., Zion G.R. Tooth retention and tooth loss in the permanent dentition of adults: United States, 1988-1991. *J Dent Res.* 1996;75: 684-95.
9. Anneloes E. Gerritsen , Dick J. Witter, Ewald M. Bronkhorst, Nico H. J. Creugers, An observational cohort study on shortened dental arches - clinical course during a period of 27–35 years. *Clinical Oral Investigations.* 2013; 17 (pp): 859-866.
10. De Oliveira BF1, Seraidarian PI2, de Oliveira SG3, Landre J Jr4, Pithon MM5, Oliveira DD6. Tooth displacement in shortened dental arches: a three-dimensional finite element study. *J Prosthet Dent.* 2014 Jun;111(6):460-5. doi: 10.1016.
11. Kourkouta S., Hemmings K.W. & Laurell L. Restoration of periodontally compromised dentitions using cross-arch bridges. Principles of perioprosthodontic patient management. *British Dental Journal* 203, 189 - 195 (2007).
12. Braun S, Bantleon HP, Hnat WP, et al. A study of bite force, part 1; Relationship to various physical characteristics. *Angel Orthod* 1995; 65(5): 367-72.
13. Waltimo A, Kononen M. A novel bite force recorder and maximal isometric bite forces values for healthy young adults. *Scand J Dent Res* 1993; 101:171–5.
14. Велески Д. Евалуација на вредностите на цвакопритисокоти реакција на потпорните ткива кај супротални протези. Докторска дисертација стом фак Скопје 1988.
15. Himmlová, Lucie & Goldmann, T & Konvickova, Svatava. (2007). MASTICATORY FORCE MEASUREMENT IN NATURAL DENTITION. *Journal of Biomechanics* 40 (2)• December 2007 with 6 Reads DOI: 10.1016/S0021-9290(07)70638-3.
16. Laurell L. Occlusal forces and chewing ability in dentitions with cross-arch bridges. *Swed Dent J Suppl.* 1985;26:160.
17. Biswas B.K.1 Bag S. 2 & Pal S.3. BIOMECHANICAL ANALYSIS OF NORMAL AND IMPLANTED TOOTH USING BITING FORCE MEASUREMENT. *International Journal of Engineering and Applied Sciences*, 2013, 4 (2): 17- 23.
18. Ye Y1, Di P1, Jia S1, Lin Y2. Occlusal force and its distribution in the position of maximum intercuspation in individual normal occlusion: a cross-sectional study. *Zhonghua Kou Qiang Yi Xue Za Zhi.* 2015 Sep;50(9):536-9.
19. Lundgren D, Laurell L. Occlusal force pattern during chewing and biting in dentitions restored with fixed bridges of cross-arch extension. I. Bilateral end abutments. *Oral Rehabil.* 1986 Jan;13(1):57-71.
20. Sato T.. STUDIES ON THE MOMENTARY OCCLUSAL FORCES OF THE NORMAL PERMANENT PREMOLARS AND MOLARS. *Nihon Hotetsu Shika Gakkai Zasshi* 1971 Volume 15 Issue 2 Pages 291-303 DOI:https://doi.org/10.2186/jjps.15.291.
21. Proeschel PA1, Morneburg T. Task-dependence of activity/ bite-force relations and its impact on estimation of chewing force from EMG. *J Dent Res.* 2002 Jul;81(7):464-8.
22. Капушевска Билјана. Функционална вредност на забите носачи на фикснопротетичките конструкции. Докторска дисертација стом фак Скопје 1998.
23. Kumagai H1, Suzuki T, Hamada T, Sondang P, Fujitani M, Nikawa H. Occlusal force distribution on the dental arch during various levels of clenching. *J Oral Rehabil.* 1999 Dec;26(12):932-5.
24. Stanišić-Sinobad D. Osnovi gnatologije. Beograd: BMG; 2001.
25. Zivko-Babić J1, Pandurić J, Jerolimov V, Mioc M, Pizeta L, Jakovac M. Bite force in subjects with complete dentition. *Coll Antropol.* 2002 Jun;26(1):293-302.
26. Sultana MH , Yamada K , and Hanada K . Changes in occlusal force and occlusal contact area after active orthodontic treatment: a pilot study using pressure-sensitive sheets. *J Oral Rehabil.* 2002;29:484–491. [Crossref] [Medline]
27. Johnsen S.E., Svensson K.G., Trulsson M., Forces applied by anterior and posterior teeth and roles of periodontal afferents during hold-and-split tasks in human subjects. *Exp Brain Res.* 2007;178(1):126-34.
28. Motta, Andréa Barreira, Pereira, Luiz Carlos. Da Cunha, Andréia R.C.C., Duda, Fernando Pereira. The Influence of the Loading Mode on the Stress Distribution on the Connector Region of Metal-ceramic and All-ceramic Fixed Partial Denture. *Artificial Organs.* 2008; 32 (4):283-291.
29. Kondo T1, Wakabayashi N. Influence of Support loss on stress and strain in premolar periodontium: a patient-specific FEM study. *J Dent.* 2009

- Jul;37(7):541-8. doi: 10.1016/j.jdent.2009.03.015. Epub 2009 Apr 19.
30. Guo Y., Tang L., Pan Y.H., [Three-dimensional finite element analysis of the stress in abutment periodontal ligament of cantilever fixed bridge under dynamic loads]. *Zhonghua Kou Qiang Yi Xue Za Zhi*. 2009 ;44(9):553-7.
 31. André Ricardo Maia Correia a João Carlos Sampaio Fernandes b José Carlos Reis Campos c Mário Augusto Pires Vaz d Nuno Viriato Marques Ramos d. Stress analysis of cantilever-fixed partial denture connector design using the finite element method. *Rev. odonto ciênc*. 2009;24(4): 420-425.
 32. Nickolay Apostolov, Ivan Chakalov, Todor Drajev. Measurement of the Maximum Bite Force in the Natural Dentition with a Gnathodynamometer. *MedInform, Journal of Medical and Dental Practic*, 2014, Vol. 1, issue 2.
 33. Jain1, Sandeep Kalra2, Vijay Prakash Mathur3, Rajath Sasidharan Pillai2 .A preliminary study to find out maximum occlusal bite force in Indian individuals. *Indian Journal of Dental Research*, Vol. 25, No. 3, May-June, 2014, pp. 325-330.
 34. Cai et al. *BMC Oral Health* (2015) 15:106 DOI 10.1186/s12903-015-0091-x.
 35. Yongqing Cai1*, Xiaoxiang Yang2 , Bingwei He2 and Jun Yao3. Finite element method analysis of the periodontal ligament in mandibular canine movement with transparent tooth correction treatment. *BMC Oral Health* (2015) 15:106 DOI 10.1186/s12903-015-0091-x.
 36. Zhou Shu-min, et al. THE STRESS ANALYSIS OF PERIOAPICAL REGION OF THE HUMAN TOOTH BY THREE-DIMENSIONAL FINITE ELEMENT METHOD http://en.cnki.com.cn/Article_en/CJFDTOTAL-BYDB198801014.htm.
 37. Jayam Bharath Kumar1, Gudimetla Jaisekhar Reddy2, Moturi Sridhar3, T Jayasimha Reddy4, Pyata Jaipal Reddy5, Sadam Srinivasa Rao6. A finite element analysis of initial stresses and displacements in the tooth and the periodontium in periodontally compromised simulations: Labial versus lingual force application. *J NTR Univ Health Sci* 2016;5:34-43
 38. FRATILAA*, C. VASILOAICAb , S. SILIVASANc, V. SEBESANd ,C. BOITORa, L.STEFa. ANALYSIS OF STRESS WITHIN THE BRIDGE AND DENTURAL PERIODONTAL AGGREGATE WITH ONE AND TWO TEETH SUPPORT USING PHOTOELASTICITY . *Digest Journal of Nanomaterials and Biostructures* Vol. 7, No. 3, July - September 2012, p. 1149 – 1155.
 39. Oruc S1, Eraslan O, Tukay HA, Atay A. Stress analysis of effects of nonrigid connectors on fixed partial dentures with pier abutments. *J Prosthet Dent*. 2008 Mar;99(3):185-92. doi: 10.1016/S0022-3913(08)60042-6.
 40. Al-Zarea B.K. Maximum Bite Force following Unilateral Fixed Prosthetic Treatment: A Within-Subject Comparison to the Dentate Side *Med Princ Pract* 2014;24:142-146. <https://doi.org/10.1159/000370214>.
 41. Панчевска С. Математички модел за определување на дистрибуцијата на оклузалните сили кај мандибуларна двострана терминална беззабност. (магистерски труд) Скопје, 2004.
 42. Stamenkovic D., Nastic M.. *Stomatoloska protetika – parcijalne proteze*. Zavod za udzbenike i nastavna sredstva, Beograd, 2000.
 43. Y. Kojima, H. Fukui. (2012) Numerical simulations of canine retraction with T-loop springs based on the updated moment-to-force ratio. *The European Journal of Orthodontics* 34:1, 10-18. Online publication date: 1-Feb-2012. [Crossref]